

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

# Urological Science

journal homepage: [www.urol-sci.com](http://www.urol-sci.com)

## CME Test

Please read this issue of Urological Science and return the postage-paid reply slip with your answers by February 28, 2013. A score of 80% or better will earn three CME credits.

- Which is the best diagnostic method for examining patients with suspected ureterolithiasis?
  - Sonography
  - KUB plain film
  - Non-enhanced helical computed tomography
  - Physical examination
  - Excretory urography
- Which one of the following descriptions of nonpalpable undescended testes is incorrect?
  - Twenty percent of undescended testes are nonpalpable
  - Five percent to 15% of nonpalpable testes are nubbing
  - Ultrasonography is necessary before surgery
  - Laparoscopy is both diagnostic and therapeutic
  - A Fowler-Stephen orchiopexy has one-step and two-step procedures

Urol Sci 2012;23(4):99-102

Urol Sci 2012;23(4):103-6

- Which of the following is the best answer of the secondary signs associated with ureteral stones on non-enhanced helical computed tomography?
  - The tissue rim sign
  - Hydroureter
  - Perirenal stranding
  - Renal density differences
  - All of the above
- Which is the best approach for the preoperative localization of an intra-abdominal testis?
  - CT scan
  - MRI
  - Ultrasonography
  - Laparoscopy
  - Angiography

Urol Sci 2012;23(4):99-102

Urol Sci 2012;23(4):103-6

Continued on next page

## Urol Sci CME Test—Reply Slip

Volume 23 Issue 4

- |        |        |        |
|--------|--------|--------|
| 1. ( ) | 2. ( ) | 3. ( ) |
| 4. ( ) | 5. ( ) | 6. ( ) |
| 7. ( ) | 8. ( ) | 9. ( ) |

Membership no.: \_\_\_\_\_

Name: \_\_\_\_\_

Score: \_\_\_\_\_

5. Which of the following is not the cause of a pear-shaped bladder?
- Pelvic lipomatosis
  - Pelvic hematoma
  - Iliopsoas muscle hypertrophy
  - Benign prostate hypertrophy

Urol Sci 2012;23(4):118-9

6. Which of the following imaging modalities is best for differentiating the causes of a pear-shaped bladder?
- Plain KUB
  - Intravenous urography
  - Sonography
  - Computed tomography

Urol Sci 2012;23(4):118-9

7. Pelvic lipomatosis occurs often in
- White males
  - White females
  - Black males
  - Black females

Urol Sci 2012;23(4):118-9

8. What is not a common urodynamic characteristic in patients with Parkinson's disease?
- Urethral sphincter pseudodyssynergia
  - Detrusor overactivity
  - Dyskinesia of the urethral sphincter
  - Underactive detrusor

Urol Sci 2012;23(4):120-3

9. What is not a characteristic of urethral sphincter pseudodyssynergia?
- High detrusor pressure during the voiding phase
  - A low maximum flow rate
  - Increased urethral sphincter activity during the voiding phase
  - A large postvoiding residual after interruption of urination
  - Low detrusor pressure during the voiding phase

Urol Sci 2012;23(4):120-3

### Volume 23 Issue 3

#### Answers:

1. (D) 2. (C) 3. (E) 4. (E) 5. (C) 6. (B) 7. (D) 8. (D) 9. (C)

廣 告 回 信	
台北郵局登記證	
台北廣字第1593號	

11051  
 台北市信義區基隆路一段432號6樓之1  
 台灣泌尿科醫學會 收